

# Mallet finger

[www.nwz.nl](http://www.nwz.nl)

# Table of contents

<b>Treatment</b>	<b>4</b>
<b>Treatment of a mallet finger with a fracture</b>	<b>4</b>
<b>Treatment of a mallet finger without a fracture (tendon injury)</b>	<b>4</b>
<b>Instructions</b>	<b>5</b>
<b>Recovery</b>	<b>5</b>
<b>Replace the splint</b>	<b>5</b>
<b>Exercises</b>	<b>5</b>
<b>Questions?</b>	<b>6</b>

**You have a mallet finger, which means that the tip of the finger lies in a bent position and you are unable to straighten it. You have snapped the tendon that normally straightens the end of the finger. The tendon may even have pulled off a small fragment of the bone.**

### Summary

- the tip of your finger lies in a bent position and you are unable to straighten it. This is called a mallet finger
- this type of injury almost always heals well without any loss of function
- a routine follow-up is not necessary for this type of injury
- the mallet finger can occur with or without a fracture, make sure you follow the treatment schedule appropriate for your injury
- you must wear the splint continuously, 24 hours a day; nonstop
- you can find more information in the folder or the Virtual Fracture Care App
- if you have any questions, or are experiencing problems with your recovery, you can call the fracture hotline 'breuklijn'  
+31 72 - 548 2465

## ■ Treatment

A mallet finger is a common injury, in most cases an operation is not necessary. You will be treated with a splint.

The treatment of a mallet finger with a loose bone fragment (fracture) is different from a mallet finger without a loose bone fragment (i.e. only tendon injury). The doctor at the emergency department will inform you about your specific type of injury.

### ■ Treatment of a mallet finger with a fracture

<b>week 0 - 6</b>	<ul style="list-style-type: none"><li>• continuously wear the splint. Do not take it off, at all</li></ul>
<b>week 6 - 8</b>	<ul style="list-style-type: none"><li>• start hand therapy (and start exercises when your therapist advises you to do so)</li><li>• wear the splint during the night and when your finger is at risk of injury</li></ul>

### ■ Treatment of a mallet finger without a fracture (tendon injury)

<b>week 0 - 8</b>	<ul style="list-style-type: none"><li>• continuously wear the splint. Do not take it off, at all</li></ul>
<b>week 8 - 10</b>	<ul style="list-style-type: none"><li>• start hand therapy (and start exercises when your therapist advises you to do this)</li><li>• wear the splint during the night and when your finger is at risk of injury</li></ul>

## ■ Instructions

Follow these instructions for a good recovery:

- you have to wear the splint at all times (24 hours a day). It is very important that you do not remove the splint. If you remove the splint before your hand-therapist advises you to do so, your recovery up to that point will be undone and the chance of a complete recovery will decrease
- if you have a tendon injury, you have to wear the splint for 8 weeks
- If you have a tendon injury with a loose bone fragment, you have to wear the splint for 6 weeks
- after these 6 weeks (bone fragment + tendon injury) or 8 weeks (isolated tendon injury), you have to wear the splint during the night for 2 more weeks. Sometimes your hand therapist advises you to wear the splint longer
- in the course of these 2 weeks, also wear the splint during the day whenever your finger is at risk of injury
- your finger and the splint should not get wet, as this will cause irritation of the skin
- are you in pain? Use a mild pain killer such as paracetamol (acetaminophen). Do not wait with the use of paracetamol until the pain is too much

## ■ Recovery

- the physiotherapist will decide when you can start exercises and will give you instructions on how to use the finger during daily activities. Be careful to gradually resume daily activities and do not make any sudden movements with your injured finger. This can damage the tendon
- most mallet injuries heal after 3 months

## ■ Replace the splint

You should never replace the splint yourself. When the splint does not fit anymore contact the fracture hotline 'breuklijn'.

## ■ Exercises

It is important that you do not start exercises before the hand therapist advises you to do this. When you can start with exercises your hand therapist will give you specific exercises.

## ■ Questions?

If you have any questions, or are experiencing problems with your recovery, you can call the fracture hotline 'breuklijn' +31 72 - 548 2465. If follow-up is necessary, we will make an appointment for you. Also contact the fracture hotline 'breuklijn' when:

- your pain increases
- the tip of your finger bends again when you have started doing the exercises
- the splint does not fit anymore
- the splint needs to be cleaned

### The Virtual Fracture Care app

Open the camera on your phone and aim it at the QR-code below. You will be automatically referred to the app.





## Noordwest Ziekenhuisgroep

Postbus 501

1800 AM Alkmaar

[www.nwz.nl](http://www.nwz.nl)

tel 072 - 548 4444

## Colofon

Redactie            chirurgie  
                          communicatie

Druk                Ricoh

Bestelnummer    313109

*Op alle onderzoeken en behandelingen van Noordwest Ziekenhuisgroep zijn de algemene voorwaarden van Noordwest Ziekenhuisgroep van toepassing, zie [www.nwz.nl](http://www.nwz.nl)*