

Greenstick fracture (child)

www.nwz.nl

Table of contents

Treatment	4
Instructions	4
Recovery	5
Reapply the brace	5
Exercises	5
Questions?	5

Your child has a small fracture in the lower-arm, which is called a greenstick fracture. This fracture is treated with a brace and a sling.

Summary

- your child has a small fracture in the lower-arm
- this type of injury almost always heals well without any loss of function
- a routine follow-up is not necessary for this type of injury
- the sling can be removed as soon as your child's pain allows this. The brace can be removed after 2 weeks. If your child is still in too much pain, the use of the brace and sling can be reduced in the third week
- to prevent the wrist from getting stiff, it is important that your child starts moving the wrist after 1-2 weeks
- you can find instructions on how to reapply the brace in the Virtual Fracture Care app
- if you have any questions, or are experiencing problems with your recovery, you can call the fracture hotline 'breuklijn':
+31 72 - 548 2465

■ Treatment

The greenstick fracture of the wrist or lower-arm is a common injury in children. This type of injury is called a greenstick fracture because it looks like a young twig that has been bent. It is treated with a brace and a sling.

week 0 - 2	<ul style="list-style-type: none">• wear the sling, remove it as soon as your child's pain allows this• wear the brace
week 2 - 3	<ul style="list-style-type: none">• wear the brace, remove it as soon as your child's pain allows
week 3 - 5	<ul style="list-style-type: none">• move the wrist and lower-arm
after 5 weeks	<ul style="list-style-type: none">• start sports

■ Instructions

Follow these instructions for a good recovery:

- the brace and sling will keep your child's arm in a comfortable position
- the sling can and should be removed as soon as your child's pain allows. Continue to wear the brace for two weeks. If your child is in too much pain after 2 weeks, you can continue to use the brace into the third week. After the third week, your child should not wear the brace anymore
- the fracture is close to the wrist. It is important that your child starts moving the wrist after 1-2 weeks. This will prevent the wrist from getting stiff and will improve healing and recovery
- after 3 weeks, your child can gradually start all usual activities again as soon as your child's pain allows this. If your child is still in too much pain, it is better to wait a few more days
- when your child takes a shower, the brace can be removed. Make sure that the arm is completely dry before putting on the brace again. You can find instructions on how to reapply the brace in the app
- is your child in pain? Use a mild pain killer such as paracetamol (acetaminophen). Do not wait until the pain is too much

■ Recovery

This type of injury almost always heals well with time and use – no specific treatment is required and routine follow-up is unnecessary.

Does the pain increase, or does it not improve? Please contact the fracture hotline 'breuklijn': +31 72 - 548 2465 .

■ Reapply the brace

If you want to reapply the brace, you can find instructions in the video in the Virtual Fracture Care app.

■ Exercises

The fracture is close to the wrist. Therefore it is important that your child starts moving the wrist after 1-2 weeks. No specific exercises are necessary.

■ Questions?

You can call us

If there are any questions after reading the information, please contact us or visit www.nwz.nl

Contact Monday to Friday between 8.30-10.00 h and 13.30-15.00 h

- breuklijn :+31 72 - 548 2465

The Virtual Fracture Care app

Open the camera on your phone and aim it at the QR-code below. You will be automatically referred to the app.



Noordwest Ziekenhuisgroep

Postbus 501

1800 AM Alkmaar

www.nwz.nl

tel 072 - 548 4444

Colofon

Redactie chirurgie
 communicatie

Druk Ricoh

Bestelnummer 313125

Op alle onderzoeken en behandelingen van Noordwest Ziekenhuisgroep zijn de algemene voorwaarden van Noordwest Ziekenhuisgroep van toepassing, zie www.nwz.nl